

Line positions and absolute intensities of nitric acid from 850 to 920 cm^{-1}
Robert A. Toth, Edward A. Cohen and Linda R. Brown, Jet Propulsion Laboratory,
California Institute of Technology, Pasadena, CA 91109

ABSTRACT

Line positions and strengths of HNO_3 from 850 to 920 cm^{-1} have been retrieved from room temperature spectra recorded at 0.00265 cm^{-1} resolution using the FTS at Kitt Peak and flowing gas samples.

SUMMARY

Some 800 transitions of ν_5 and 550 lines of $2\nu_9$ have been measured with precisions of 3 to 8%. Measurements of an additional 438 features with ambiguous assignments have also been obtained with 10% - 15% precisions. These include a few strong absorptions between 885.414 cm^{-1} and 885.470 cm^{-1} which are recognized as the Q-branch of $\nu_5 + \nu_9 - \nu_9$. The measured line intensities and band strengths of the ν_5 and $2\nu_9$ bands have been compared to prior measurements and to values given in the HITRAN compilation; the average of the ratios, this work/HITRAN, was found to be $0.92 \pm 11\%$. The analysis of these data has been greatly hindered by the high density of transitions and the lack of knowledge about several other hot bands in the region.

The research described in this paper was carried out by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.